

Parasitism potential of *Diadegma insulare* (Cameron) (Ichneumonidae: Hymenoptera) against *Plutella xylostella* (L.) (Plutellidae: Lepidoptera)

AM Bhosale^{1*}, VA Bhosale²

^{1,2} Department of Agrochemicals and Pest Management, Shivaji University, Kolhapur, Maharashtra, India

Abstract

Diadegma insulare (Cameron) (Ichneumonidae: Hymenoptera) is a solitary larval parasitoid of *Plutella xylostella* (L.) (Plutellidae: Lepidoptera), a notorious and polyphagous pest of cabbage and cauliflower in India. Nutritional requirement, host age, specificity and density dependent factor play a very crucial role in biocontrol program of parasitoids and further their release in the field for pest control strategies. Attempt has been made to initiate their mass multiplication for successful biocontrol programme. Therefore, present work was conducted to find out the optimum host stage and density for maximum progeny production of the parasitoid under laboratory conditions. The parasitoid caused highest mortality in the pest larvae of second instars, 4 day old larvae were attacked most with high percent parasitism, 39.00%. Optimum density for maximum progeny production of *D. insulare* was 20, which generate maximum parasitism (41.00%). Host specificity by exposing the parasitoids towards different host species and analyse parasitoid preference by *P. xylostella* > *Spodoptera litura* (Fabricius) > *Mythimna separate* Walker > *Achaea janata* (Linnaeus). Nutritional requirement of parasitoid was tested with different foodstuffs and found 50% honey best suited for maximum longevity 8.2 and 11.4 days for males and females respectively. The longevity ratio also female biased, 1: 1.39 (Male: Female). From the results it concludes that *D. insulare* fed with 50% honey solution, exposed to 3-5 day old caterpillars of *P. xylostella* at density of 20 gave maximum progeny production and effectively utilized in the biocontrol programme.

Keywords: *Diadegma insulare*, parasitism potential, biocontrol, parasitoids, *Plutella xylostella*, pest management

Introduction

Agriculture is the basic component of Indian economy. Development and use of improved practices enhance the status of farmers as well as their community. Farmers are facing problem of pests and diseases of crop plants. Long ago the pests and diseases could be curtailed environmentally but then it shifted to era of chemical control. Due to residual problems, cost of sprays, lack of labours, resistance problem, secondary pest outbreak, pest resurgence and phytotoxicity of insecticides the farmers now deliberately moved towards biological control of agricultural pests. Biological control of different pests with biocontrol agents enhance the crop yield and also improve the quality of produce with least residual problem. The above fact clearly indicates that there is extreme need of minimizing pesticidal use. Therefore, attempts have been made with the use of biocontrol agents for effective control of pests as ecofriendly control measure.

Plutella xylostella (L.) (Plutellidae: Lepidoptera) is the most notorious pest of cabbage and cauliflower crops in India. Several management practices were implemented to achieve the control of the pest. Among biocontrol agents *Diadegma insulare* (Cameron) (Ichneumonidae: Hymenoptera) is most common and potent solitary larval parasitoid scattered in nature and control the pest population effectively. Host searching and selection of host density by parasitoid counts the success of biocontrol programme of any pest species (Sathe and Bhosale, 2011) [15]. The high rate of parasitism is desirable character of an ideal parasitoid. In colonization and mass production of

parasitoids in biocontrol strategies the shape, size, nutritional suitability and age of host plays important role (Vinson, 1976; Vinson and Iwantsch, 1980).

Leong and Oatman (1968) [8], Lewis and Vinson (1971) [9], Lingren and Nobel (1972) [11], Pawar *et al.* (1989) [13], Romeis and Shanower, (1996) [14], King (1998) [7], Wackers (2001) [19], Eliopoulos (2007) [4], Khatri *et al.* (2012) [6], Sathe *et al.* (2012) [15], Bhosale (2018) [1] were made investigations on optimum age, density and specificity of hosts and nutritional requirement of ichneumon parasitoids. The present study was carried with *D. insulare*, a larval parasitoid of *P. xylostella* to find out the optimum age of host for obtaining maximum progeny of parasitoids, which will help in mass rearing and field release for effective biocontrol program.

Materials and methods

Rearing of host species

P. xylostella larvae were reared in small perforated plastic container (7x8 cm, Diameter x Height). After adult emergence they may transferred in oviposition cage 25x25x25 cm (LxWxH). First instar caterpillars usually hatch after 2 days from oviposition. These larvae were collected and used for further experiment. During the course of study, the host caterpillars were fed with cabbage leaves. Similarly, the hosts used to conduct the host specificity experiment were reared their natural food like, *Spodoptera litura* (Fabricius) and *Achaea janata* (Linnaeus) on leaves of castor *Ricinus communis* L. and *Mythimna separate* Walker on leaves of maize *Zea mays* L.

Rearing of parasitoid

Adults of *D. insulare* were reared in ventilated wooden cage (30x30x30 cm, LxWxH). *D. insulare* are very small insects and are negatively geotropic, hence cages must be made with glass walls on three sides and top of the cage while one wall was made up of very fine mesh cloth for proper handling of parasitoids. The *D. insulare* was fed with 50% honey solution. Adults of parasitoids released for oviposition in the rearing cages for 24 h with different age and densities of *P. xylostella* caterpillars. After 24 h, adults were removed and hosts reared for further analysis. The cocoons of parasitoids then transfer into separate container and adults of *D. insulare* emerges out that can be used for experimental purpose.

Nutritional requirements and adult longevity of parasitoid on different foodstuffs

Emerged adults of *D. insulare* are fed with different foodstuffs to analyze the ideal feed for getting highest longevity. Honey acts as natural food for any parasitoid, hence three concentrations of 100, 50 and 10 percent were made to analyze the longevity of parasitoid. However, 50 percent glucose and sucrose; and juice of citrus and apple can also provide as a food.

Pest age related parasitism

To determine the effect of host age on parasitism, 20 larvae of *P. xylostella* of known age (ranging from less than 1 day to 13 days old) were exposed to single mated female of *D. insulare* in a glass cage for 24 hrs. The larvae were removed and placed in separate containers for further observations. Daily records of cocoon construction and parasitoid emergence from each container were neatly observed.

Host density for optimum parasitization

P. xylostella caterpillars (4-5 day old) were exposed in densities of 10, 20, 30, and 40 towards mated females of *D. insulare* for 24 hrs in oviposition cage 25x25x25 cm (LxWxH). The host larvae were reared into plastic containers to record further development or parasitoid emergence.


Host specificity for optimum parasitization

Host specificity was conducted by exposing the mated females of parasitoid towards caterpillars of different host species like *P. xylostella*, *S. litura*, *M. separata* and *A. janata*. The hosts were placed in the oviposition cage for 24 h. Hosts were released in 20 densities to record optimum parasitism. Afterwards the hosts were reared on the natural diet and observe the emergence of parasitoid or further lifecycle of host species.

The experiments were carried out at 25±2°C, 60±5% RH and 12hr. photoperiod. During the course, cabbage leaves were provided as a food to the caterpillars of *P. xylostella* and other appropriate food for other experimented host species, while the parasitoids were fed with 50% honey solution. Each experiment was repeated 5 times for confirming the result.

Results and Discussion


D. insulare was most effective in controlling the caterpillars of *P. xylostella* than other biocontrol agents associated with particular pest. The results of host specificity experiment (Fig. 1) revealed that the parasitoid prefer *P. xylostella* as the primary host with 40.00% parasitism. Among tested hosts, parasitoid showed 19.00 percent parasitism for *S. litura*, 15.00 percent parasitism for *M. separata* and 6.00 percent parasitism for *A. janata*. The order of preference for parasitism shown by the parasitoid was *P. xylostella* > *S. litura* > *M. separata* > *A. janata*.



*Each value is the mean of five replicates with error bars indicating standard error of mean (SEM).

Fig 1: Host specificity of *D. insulare*

Adult longevity of *D. insulare* with different foodstuff were analyzed and plotted in fig. 2. Parasitoid survived longer with 50% honey solution with maximum male: female longevity ratio (1: 1.39). The maximum longevity of male and female when fed with 50% honey solution was 8.2 and 11.4 days respectively. Hence, it could be best suited for mass rearing of parasitoid in the laboratory.



*Each value is the mean of five replicates with error bars indicating standard error of mean (SEM).

Fig 2: Adult longevity of *D. insulare* with different foodstuffs

The results recorded in table 1 indicate that the parasitoid caused highest mortality in the second instar caterpillars. The caterpillars of 3-6 days old are preferred for parasitism whereas, beyond 11 days old were not attacked by the parasitoid. Four day old caterpillars were attacked most with high percent parasitism (39.00%).

Table 1: Host age related parasitism by *D. insulare*

Host age (days)	% Parasitism	% Mortality	% Moth emergence
1	4.00 (±2.20) ^{ef}	6.00 (±3.70) ^a	90.00 (±2.50) ^{ab}
2	8.00 (±3.70) ^{de}	8.00 (±1.20) ^a	84.00 (±3.70) ^{abcd}
3	22.00 (±3.70) ^{abc}	10.00 (±1.00) ^a	68.00 (±3.00) ^{de}
4	39.00 (±2.50) ^a	8.00 (±2.90) ^a	53.00 (±1.90) ^e
5	25.00 (±3.70) ^{ab}	9.00 (±2.50) ^a	66.00 (±4.60) ^{de}
6	20.00 (±3.40) ^{abcd}	9.00 (±1.20) ^a	71.00 (±2.70) ^{cde}
7	16.00 (±0.00) ^{bcd}	11.00 (±5.10) ^a	73.00 (±5.10) ^{bcd}
8	9.00 (±1.90) ^{cde}	4.00 (±3.40) ^a	87.00 (±1.90) ^{abcd}
9	9.00 (±1.90) ^{cde}	10.00 (±3.30) ^a	81.00 (±4.60) ^{abcd}
10	4.00 (±1.20) ^{ef}	7.00 (±3.30) ^a	89.00 (±2.40) ^a
11	3.00 (±2.00) ^{ef}	10.00 (±2.70) ^a	87.00 (±2.50) ^{abc}
12	0.00 (±0.00) ^f	11.00 (±4.00) ^a	89.00 (±4.00) ^a
13	0.00 (±0.00) ^f	10.00 (±3.40) ^a	90.00 (±3.40) ^{ab}
CD (P=0.05)	11.57	17.36	15.63

*The data presented are the mean of five replicates. Different letters indicate the significant difference (One way ANOVA) P<0.05 Tukey's standardized range (HSD) test. Figures in parentheses are standard error of mean (SEM).

The results of optimum host density for maximum progeny production of parasitoid were tabulated in table 2 showed that the number of parasitoids obtained from host density 20 was highest with 41.00 percent parasitism, compared to those produced from other host densities 10, 30 and 40 with 16.00, 32.67, 30.00 and 24.80 mean percentage of parasitism respectively.

Table 2: Host density dependent parasitism by *D. insulare*

Host density	% parasitism	% Mortality	% Moth emergence
10	19.00 (±1.61) ^b	20.00 (±0.24) ^a	61.00 (±1.61) ^a
20	41.00 (±1.12) ^a	18.00 (±1.47) ^a	40.00 (±2.38) ^b
30	32.67 (±0.84) ^a	14.33 (±0.61) ^a	53.00 (±1.14) ^{ab}
40	29.00 (±0.87) ^{ab}	17.50 (±0.30) ^a	53.50 (±1.16) ^{ab}
50	23.80 (±0.43) ^{ab}	14.60 (±0.37) ^a	61.60 (±0.64) ^{ab}
CD (P=0.05)	11.60	7.07	13.68

*The data presented are the mean of five replicates. Different letters indicate the significant difference (One way ANOVA) P<0.05 Tukey's standardized range (HSD) test. Figures in parentheses are standard error of mean (SEM).

The statistical analysis was made by one-way ANOVA using the statistical software package SAS 9.3(32) English. The percent values were transformed to arcsine values before analysis.

Sathe and Bhosale (2011) [15] reported the host density 100 for obtaining maximum progeny production (38.50%) of the parasitoid *D. insulare*. In present findings 20 host density shows maximum parasitism (41.00%). Likely, Sathe *et al.* (2012) [16], found the optimum host density for maximum parasitism of *Apanteles cretonoti* Viereck on *Amsacta moorei* Butler which was 20 with parasitism of 50 percent. Pawar *et al.* (1989) [13], studied the parasitism of *Campoletis chlorideae* Uchida on *Helicoverpa armigera* (Hubner), they found average percentage parasitism of first to third instar larvae, which are only parasitised by parasitoid, parasitism found on associated crop was 44.2 on sorghum, 33.1 on chickpea, 32.6 on pearl millet, 7.1 on groundnut and 4.2 on pigeonpea.

Lingren *et al.* (1970) [10] stated the host age preference of *C. chlorideae* towards four lepidopterous host species *Prodenia ridinia* (Craner), *Prodenia praefica* Grote, *Trichopulsia ni* (Hubner) and *Pseudoletia unipuncta*

(Hawarth). They reported that caterpillars 1-8 day old of all hosts were susceptible for parasitism, 2-6 day old being most acceptable. In present findings 2-9 day old caterpillars of *H. armigera* were susceptible, 3-7 day old caterpillars readily accepted and 4-5 day old being most suitable for parasitism.

Nikam and Basarkar (1981) [12] studied the reproductive potential of *C. chlorideae* and reported maximum parasitization at host density 40. In *Campoplex haywardi* Blanchard, an ichneumonid parasitoid of *Phthorimaea operculella* Zeller, the optimum host density was 75 larvae per tuber for maximum progeny production (Leong and Oatman, 1968) [8]. Similarly, Cardona and Oatman (1971) [2], reported 90 host density of *Keiferia lycopersicella* (Walsingham) as optimum number for maximum parasitism by *Pseudapanteles (=Apanteles) dignus* Muesebeck. In *P. dignus*, they reported the percentage of parasitization increased with the increase in number of hosts (30, 60 and 90) up to host density 90 per replicate. A decrease in parasitization observed in all replicates when 120 larvae were offered. In present study percent parasitism was found decreasing beyond 20 host density that suggesting the suitability of the larval number.

Han *et al.* (2013) [5] studied the host preference and suitability in *C. chlorideae* and recorded the parasitism against hosts *H. armigera*, *M. separata* and *Spodoptera exigua* (Hubner). They found parasitoid showed maximum parasitism on *H. armigera* followed by *M. separata* and *S. exigua*. In present findings the parasitoid preferred *P. xylostella* later *S. litura*, *M. separata* and then *A. janata*. Dhillon and Sharma (2007) [3], recorded survival and development of *C. chlorideae* on various insect and crop hosts and found maximum cocoon formation (82.4%) and adult emergence (70.5%) on *H. armigera* than other hosts. Their studies focused on host preference behaviour of *C. chlorideae* on alternate insect hosts like *S. exigua*, *M. separata* and *A. janata*. They noted that the maximum parasitism found on *H. armigera*, followed by *M. separata*, *S. exigua* and *A. janata*. In present findings highest parasitism noted on *P. xylostella* with *S. litura* having second position followed by *M. separata* and *A. janata*.

D. insulare has been successfully initiating the biocontrol program for managing the *P. xylostella*. Parasitoid can be mass reared in laboratory scale on 50% honey solution. For getting maximum progeny of parasitoid exposed towards second instar *P. xylostella* caterpillars with 20 host density. The mass rearing of parasitoid may initiate the biocontrol programme for *P. xylostella*.

Acknowledgments

Author is thankful to Coordinator, Department of Agrochemicals and Pest Management, Shivaji University, Kolhapur (Maharashtra) for providing necessary facilities and Golden Jubilee Research Fellowship, Shivaji University, Kolhapur (Maharashtra) for providing financial assistance to research project.

References

- Bhosale AM. Parasitism potential of *Campoletis chlorideae* Uchida (Hymenoptera: Ichneumonidae) against *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae). Entomon. 2018; 43(3):171-176.

2. Cardona C, Oatman ER. Biology of *Apanteles dignus* (Hymenoptera : Braconidae), a Primary parasite of the tomato pin worm. *Ann. Ent. Soc. Am.* 1971; 5:996-1007.
3. Dhillon MK, Sharma HC. Survival and development of *Campoletis chlorideae* on various insect and crop hosts: implications for Bt-transgenic crops. *Journal of Applied Entomology.* 2007; 131(3):179-185.
4. Eliopoulos PA. The importance of food supplements for parasitoids of stored product pests: the case of *Venturia canescens* (Hymenoptera: Ichneumonidae). *Integrated Protection of Stored Products.* 2007; 30(2):37-41.
5. Han LB, Huang LQ, Wang CZ. Host preference and suitability in the endoparasitoid *Campoletis chlorideae* is associated with its ability to suppress host immune responses. *Ecological Entomology.* 2013; 38:173-182.
6. Khatri D, He XZ, Wang Q. Reproductive fitness of *Diadegma semiclausum* Hellen (Hymenoptera: Ichneumonidae) in response to host density. *New Zealand Plant Protection.* 2012; 65:142-147.
7. King BH. Host age response in the parasitoid wasp *Spalangia cameroni* (Hymenoptera: Pteromalidae). *Journal of Insect Behavior.* 1998; 11(1):103-117.
8. Leong GKL, Oatman ER. The biology of *Campoplex haywardi* (Hymenoptera: Ichneumonidae), a primary parasite of the potato tuber worm. *Ann. Ent. Soc. Am.* 1968; 61:26-36.
9. Lewis WJ, Vinson SB. Suitability of certain *Heliothis* (Lepidoptera: Noctuidae) hosts for the parasite, *Cardiochiles nigriceps*. *Ann. ent. Soc. Am.* 1971; 64:970-972.
10. Lingren PD, Guerra RJ, Nickelsen JW, White C. Hosts and host-age Preference of *Campoletis perdistinctus*. *Journal of Economic Entomology.* 1970; 63(2):518-522.
11. Lingren PD, Nobel LW. Preference of *Campoletis perdistinctus* for certain noctuid larvae. *J. Econ. Entomol.* 1972; 65:104-107.
12. Nikam PK, Basarkar CD. Influence of host density on reproductive potential of *Campoletis chlorideae* (Uchida) (Hymenoptera: Ichneumonidae), an internal parasite of *Heliothis armigera* (Hubn.). *Bioresearch.* 1981; 5:101-104.
13. Pawar CS, Bhatnagar VS, Jadhav DR. *Campoletis chlorideae* Uchida (Hymenoptera: Ichneumonidae) as a parasite of *Helicoverpa armigera* (Hub.) (Lepidoptera: Noctuidae) in southwest India. *Proc. Animal Sciences.* 1989; 98(4):259-265.
14. Romeis J, Shanower TG. Arthropod natural enemies of *Helicoverpa* (Hubner) (Lepidoptera: Noctuidae) in India. *Biocontr Sci Tech.* 1996; 6:481.
15. Sathe TV, Bhosale AM. *Plutella xylostella* (L.) density requirement for maximum progeny production of *Diadegma insulare* (Cameron). *Asian J. Animal Sci.* 2011; 6(2):212-214.
16. Sathe TV, Bhosale AM, Jadhav AD, Patil RN. Host density for optimum parasitization by *Apanteles creatonoti* Viereck (Braconidae: Hymenoptera) on *Amsacta moorei* Butler (Arctiidae: Lepidoptera). *J. Appl. Zool. Res.* 2012; 3(1):19-22.
17. Vinson SB. Host selection by insect parasitoids. *Annual Review of Entomology.* 1976; 21:109-133.
18. Vinson SB, Iwantsch GB. Host suitability for insect parasitoids. *A. Rev. Ent.* 1980; 25:397-419.
19. Wackers FL. A comparison of nectar and honeydew sugars with respect to their utilization by the hymenopteran parasitoid *Cotesia glomerata*. *Journal of Insect Physiology.* 2001; 47:1077-1084.