



The effectiveness of biological mosquito larvicide: Bacteria (*Bacillus thuringiensis israelensis*) embedded in Silica Granules

Milagros M Greif

Department of Biology, College of Arts and Sciences Cebu Normal University Osmeña Boulevard, Cebu, Philippines

Abstract

Mosquitoes are very tiny insects but the impact they brought to humans and society is very huge. An application technique for the control of mosquito larvae under tropical conditions was conducted in Cebu City, Philippines. The study used a bacterium, *Bacillus thuringiensis israelensis* embedded in silica granules under laboratory conditions. Five concentrations were prepared in range finding and 4 concentrations in definitive range finding tests. Each of the concentration was tested with 25 3rd and early 4th instar larvae of *Aedes aegypti*. Results obtained from the test have very effective residual effects against the larvae of mosquitoes. The efficacy varies according to the concentrations applied to the mosquito larvae. The high mortality rate of mosquito larvae is associated numerically with higher concentration of the larvicide (1.6 grams) and the low mortality rate of mosquito larvae is associated numerically with lower concentration (0.4 grams) after one (1) hour application of the larvicide. This means that even, with lower level of product concentration of the larvicide, the mortality rates among mosquito larvae are still high with the lowest rate of 93% after 1 hour of exposure to the biological mosquito larvicide. Further, there is a significant increase in the number of dead mosquito larvae after 2 hours of treatment. It implies that the toxicity level of the larvicide is still there even after 2 hours of treatment. However, 1.2 g concentration is proven to be effective and achieved 100% mortality after 2 hours of application to the mosquito larvae.

Keywords: *Aedes aegypti*, *Bacillus thuringiensis israelensis*, silica granules, mosquito, biological control, dengue control

Introduction

Mosquitoes are very tiny insects but the impact they brought to humans and society is very huge. They are carriers of diseases like dengue and malaria that can kill humans. Dengue is the world's most important mosquito-borne virus disease, with 2500 million people worldwide at risk of infection and 20 million cases a year in more than 100 countries [1]. It is transmitted by the bite of a female *Aedes* mosquito. In the Philippines, dengue is a high-profile and ongoing public health concern together with the newly emerged zika virus. The Philippines experienced the seventh highest number of dengue cases in the world between 2004 and 2010 [2]. There are 179,540 cases that were infected on the virus from January to November 2018 with 907 deaths [3]. In Central Visayas, there are 15,171 dengue cases with the highest infection cases of 2,660 resulting to 14 deaths in Cebu City from January 2018- November 2018 [4]. Although there is already dengue vaccine available in the market however, it poses controversial issues in the society. Due to this, one way of reducing the risk of being infected by mosquito borne diseases is to control the mosquito while they are still in the water or during larval stages. Controlling mosquito larvae in the water is either by the application of Chemical (ex. temephos) or biological way (bacterium). Unfortunately, the widespread use of chemical insecticides has contributed to increasing resistance to these agents among *Aedes aegypti*, especially in the Americas and the Caribbean [5]. Further, one of the disadvantages of chemical application is that there are also other organisms that are affected upon the application. Published international scientific studies found out that temephos is toxic to some bird species and other organisms. Due to this, it cannot be considered as environmentally friendly. A very promising

and really environmentally friendly tool to fight against dengue, which is used internationally and highly recommended by World Health Organization (WHO), is the bacterium *Bacillus thuringiensis israelensis* (B.t.i.). This bacterium is proven to be only toxic to the target enemies, namely: larvae of mosquitoes, black flies and midges (responsible for human diseases) based on international scientific researches and publications. The study evaluates the effectiveness of biological mosquito larvicide *Bacillus thuringiensis israelensis* in liquid formulation as formulated by the author and registered by Cebu Normal University as utility model by the Bureau of Patents, Philippines. Specifically it aims to determine: the mortality rate of mosquito larvae by the application of four concentrations of the larvicide in liquid formulation and; the efficacy of the larvicide after 24 hour of treatment to the mosquito larvae.

Materials and Methods

Mosquito Larvae Collection

Mosquito larvae were collected from clean & stagnant mosquito breeding sites using a water dipper. The collected larvae were kept in half closed jar bottles & transported to the Biology Laboratory, Cebu Normal University. In the laboratory, the mosquito larvae samples were poured into the plastic basin. The larvae in the basin were sorted according to the ages and only the 3rd & early 4th instar larvae of *Aedes mosquito* were used for the experiment.

Range Finding Test of Utility Model Patent (Assessment of the potency and the minimum effective dosage in the laboratory)

The prototype of the product was provided by the researcher which is the maker of the utility model. The utility model on

the process of producing mosquito larvae control comprising the steps of:

1. mixing 1 to 100g of *Bacillus thuringiensis israelensis* with 1 liter of tap water in a mixing vessel;
2. putting 5 kilograms silicate granules in a container;
3. pouring the mixture of *Bacillus thuringiensis israelensis* and tap water in the container containing the silicate granules; and
4. Stirring thoroughly until the mixture will be absorbed by the silicate granules.

Samples of mosquito larvae were collected in Cebu City. Twenty five mosquito larvae samples were used per concentration and per concentration had triplicates. There were 5 concentrations of the product that were tested under laboratory conditions on the mosquito larvae, namely: 2 g, 4 g, 6 g, 8 g and 10 g. Twenty five 3rd & early 4th instar larvae of *Aedes aegypti* (Linn.) in 2 ml of water were added to plastic disposable cups filled with 148 ml of distilled water. Mortality readings of larvae were taken in five readings after the application of the larvicide (after 1 hr, 2, 4, 8, & 24 hrs). If the desired mortality rate (100%) was not attained, another batch of mosquito larvae was released to higher concentration until a 100% mortality rate of the larvae was achieved.

Definitive Test of Utility Model Patent

Four different concentrations based on the results from range finding test were tested namely: 0.4g, 0.8g, 1.2g, & 1.6g in triplicates. The same procedure in the range finding test was applied for the application of the different concentrations.

Data Analysis

Data collected on mortality rate difference of mosquito larvae among the four concentrations were calculated using ANOVA.

Results and Discussion

Range Finding Test of Utility Model Patent (Assessment of the potency and the minimum effective dosage in the laboratory)

Results showed that after 1 Hr of the application of the product, average mortality rate of 100% was attained in 2,4,6,8, & 10 grams concentrations. Based on the results, 2 g concentration was selected for definitive test considering that after 2 hrs. of the product application, the lowest concentration (2g) was able to attain a 100% mortality rate of mosquito larvae. Figure 1 showed the results of the Range Finding Test.




Fig 1: Percentage Mortality Rate of Mosquito Larvae in Silica Range Finding Test.

Definitive Test of Utility Model Patent

The product showed a very promising result as shown in Figure 2. After 1 hr of product application, 93%, 96%, 97%, & 99% mortality rates of mosquito larvae in 0.4g, 0.8g, 1.2g, & 1.6g concentrations were attained, respectively. Further after 2 hrs. of product application, mortality rate of 99% in 0.4 & 0.8 g concentrations and 100% mortality rate was attained in 1.2 & 1.6 g concentrations.




Fig 2: Percentage Mortality Rate of Mosquito Larvae in Silica Definitive Test.

The descriptive statistics associated with the mortality rate of mosquito larvae after the application of four different concentrations of biological mosquito larvicide embedded in silica granules are reported in Table 1 below. It can be seen that higher mortality rate of mosquito larvae ($M = 24.7, SD = 0.58$) was associated numerically with higher larvicide concentration embedded in silica granules (1.6 grams) and the low mortality rate of mosquito larvae ($M = 23.3, SD = 0.58$) was associated numerically with lower concentration (0.4 grams) after one (1) hour application of the larvicide.

Table 1: Descriptive Statistics of Mortality Rates of Mosquito Larvae across Four Different Concentrations of Biological Mosquito Larvicide in Silica Granules

Concentration	N	M	SD	Mean Mortality Percentage	Levene's Statistic	P - value
0.4 grams	25	23.3	0.58	93.2%	F(3,8) = 0.333	.802
0.8 grams	25	24.0	1.00	96.0%		
1.2 grams	25	24.3	0.58	97.2%		
1.6 grams	25	24.7	0.58	98.8%		

Prior to conducting the ANOVA analysis, the assumption of normality was evaluated and determined to be satisfied as the group's distributions were associated with skewness and kurtosis less than | | and | |⁶; Table 1). Moreover, the assumption of homogeneity of variances was tested and satisfied based on Levene's F-test, $F(3, 8) = 0.333, p = .802$. In order to test the hypothesis that the level of concentration of biological mosquito larvicide in silica granules had an effect on mortality rate of mosquito larvae, one-way between-Subjects ANOVA was performed as displayed in Table 2. The independent between-groups ANOVA yield a statistically non-significant effect, $F(3, 8) = 1.944, p = .201$. Thus, the null hypothesis of no differences in the mean mortality rates of mosquito larvae when exposed to different larvicide concentrations embedded in silica granules was not rejected. This means that even with lower level of larvicide concentration, the mortality rates among mosquito larvae are still high with the lowest rate of 93.2% after 1 hour of

exposure to the biological mosquito larvicide. However, a moderate 42.2% variance in the mortality rates of mosquito larvae was accounted for by the larvicide concentration as indicated by its effect size.

Table 2: ANOVA Summary Table of Different Levels of Larvicide Concentration in Silica Granules towards the Mortality Rates of Mosquito Larvae after 1 Hour.

Variation Squares	DF	Mean Square	F - value	P - value
Between Groups 2.917	3	0.972		
Within 4.000	8	1.944	2.01	0.500
Total 6.917	11			

Furthermore, another one-way between-subjects ANOVA was performed after 2 hours of larvicide application to the mosquito larvae as displayed in Table 3. The independent between-groups ANOVA yield a statistically non-significant effect, $F(3, 8) = 0.667, p = .596$, thus, the null hypothesis of no differences in the mean mortality rates of mosquito larvae after 2 hours exposure to different larvicide concentrations in silica granules was not rejected. This means that even, with lower level of larvicide concentration, the mortality rates among mosquito larvae are still high with the lowest rate of 98.7% to the highest rate of 100% after 2 hours of exposure to the biological mosquito larvicide. However, a noticeable 20.0% variance in the mortality rates of mosquito larvae was accounted for by the larvicide

concentration as indicated by its effect size.

Table 3: ANOVA Summary Table of Different Levels of Larvicide Concentration in Silica Granules towards the Mortality Rates of Mosquito Larvae after 2 Hours

Sources of Variation	Sum of Squares	DF	Mean Square	F - value	P - value
Between Groups	0.333	3	0.111		
Within	1.333	8	0.667	5.96	0.167
Total	1.667	11			

In order to test the hypothesis that the mortality rates of mosquito larvae is different in different time intervals after larvicide treatment in silica granules, a paired *t*-test was conducted as displayed in Table 4 below.

An alpha level of 0.05 was still utilized in this analysis. Descriptive statistics show that there were no violations of normality and homogeneity of variances in the data gathered. A statistically significant difference was evident in the mortality rates of mosquito larvae between 1 hour ($M = 96.3\%, SD = 3.2\%$) and 2 hours after ($M = 99.3\%, SD = 1.6\%$) larvicide treatment, $t(11) = 3.00, p = .012$. This simply means that there is a significant increase in the number of dead mosquito larvae after 2 hours of treatment. It implies that the toxicity level of the larvicide in silica granules is still there even after 2 hours of treatment. A large Cohen's effect size was noted in this study, $d = 0.866$.

Table 4: Differences in the Mortality Rates of Mosquito Larvae between 1 hour and 2 hours after Treatment of larvicide in Silica Granules.

No. of hours After Treatment	N	M	SD	Mean Percentage Difference	t - value	P - value
After	12	96.3%	3.2%	3.0%	3.00	.012
1 hour	12	99.3%	1.6%			
After 2 hours						

The results obtained from the test have very high efficacy against the larvae of mosquitoes. The efficacy depends on the concentrations applied to the mosquito larvae. As shown in the results, high mortality rate of mosquito larvae is associated numerically with higher concentration of the larvicide (1.6 grams) and the low mortality rate of mosquito larvae is associated numerically with lower concentration (0.4 grams) after one (1) hour application of the larvicide.

This means that even, with lower concentration of the larvicide, the mortality rates among mosquito larvae are still high with the lowest rate of 93% after 1 hour of exposure to the biological mosquito larvicide. Further, there is a significant increase in the number of dead mosquito larvae after 2 hours of treatment. It implies that the toxicity level of the larvicide is still there even after 2 hours of treatment. However, 1.2 g concentration is proven to be effective and achieved the same results compared with higher concentrations after 2 hours of application to the mosquito larvae. The larvicidal efficacy is due to the parasporal crystal of *B. t. i.* that composed of four major proteins namely: 27, 65, 128 and 135 kDa [7]. The high efficacy that Bti showed in laboratory and field trials during the early 1980s led rapidly to its development as a commercial bacterial larvicide for control of mosquito and blackfly larvae [8, 9].

Laboratory studies suggest that *B.t.i.* did not show any resistance to mosquito larvae due primarily to the presence of Cyt1A in the parasporal body [10, 11]. Cyt1A's capacity to synergize endotoxin proteins, including the *B. sphaericus* Bin toxin against resistant and non-sensitive mosquitoes [12].

Conclusion

The product used in the study that is registered by Cebu Normal University as utility model by the Bureau of Patents is a very promising tool to combat the problem of *Aedes* mosquitoes which are the carrier of dengue and other viruses.

Acknowledgements

The author wishes to thank Mrs. Estrellita Capulan, biology laboratory custodian for the assistance in the collection of mosquito larvae samples. This research work was partially funded by Cebu Normal University.

References

- World Health Organization. 2017. www.who.int/news-room/fact-sheets/detail/dengue-and-severe-dengue
- World Health Organization. 2015. Vector Borne Diseases.
- Department of Health
- (DOH-7), Regional Epidemiology Surveillance Unit (RESU). 2018.
- Department of Health (DOH-7), Regional Epidemiology Surveillance Unit (RESU), 2019.
- Polson KA, Brogdon WG, Rawlins SC, Chadee DD. Characterization of insecticide resistance in Trinidadian strains of *Aedes aegypti* mosquitoes. *Acta Trop*, 2010. doi:10.1016/j.actatropica.2010.09.005. Google Scholar.
- Schmider E, Ziegler M, Danay E, Beyer L, Bühner M. Is it really robust? Reinvestigating the robustness of ANOVA against violations of the normal distribution

- assumption. *Methodology*,2010:6:147-151. [Google Scholar]
8. Crickmore N, Zeigler DJ, Feitelson J, Schnepf E, Lambert B, Lereclus D *et al.* Revision of the Nomenclature for the *Bacillus thuringiensis* Pesticidal cry Genes. Program and Abstracts of the 28th Annual Meeting of the Society for Invertebrate Pathology, 1995.
 9. Mulla MS. Activity, field efficacy, and use of *Bacillus thuringiensis israelensis* against mosquitoes. In *Bacterial Control of Mosquitoes and Black Flies* (ed. H. de Barjac and D. J. Sutherland), 1990, 134-160. New Brunswick: Rutgers University Press. Google Scholar.
 10. Becker N, Margalit J. Use of *Bacillus thuringiensis israelensis* against mosquitoes and blackflies. In *Bacillus thuringiensis, An Environmental Biopesticide: Theory and Practice* (ed. P. F. Entwistle, J. S. Cory, M. J. Bailey and B. J. Higgs), 1993 147-170. New York: John Wiley & Sons. Google Scholar.
 11. Georghiou GP, Wirth MC. Influence of exposure to single versus multiple toxins of *Bacillus thuringiensis* subsp. *israelensis* on development of resistance in the mosquito *Culex quinquefasciatus* (Diptera: Culicidae). *Appl. Environ. Microbiol*,1997:63:1095-1101. Google Scholar.
 12. Wirth MC, Georghiou GP, Federici BA. CytA enables CryIV endotoxins of *Bacillus thuringiensis* to overcome high levels of CryIV resistance in the mosquito *Culex quinquefasciatus*. *Proc. Natl. Acad. Sci. USA* **94**,10536-10540. Google Scholar, 1997.
 13. Wirth MC, Walton WE, Federici BA. Cyt1A from *Bacillus thuringiensis* restores toxicity of *Bacillus sphaericus* against *Culex quinquefasciatus* (Diptera: Culicidae). *J. Med. Entomol*, 2000a:37:401-07. Google Scholar.