



Proximate analysis of *Labeo rohita* specific growth rate (SGR) and fingerlings exposed to titanium dioxide (TiO₂ NPs) of Bi-modal effect of growth

Gohila B*, Damodaran R, Azhagu Madhavan S

PG & Research Department of Zoology and Biotechnology, A Veeriyar Vandayar Memorial Sri Pushpam College, (Autonomous) Poondi, Thanjavur (Dt), Tamil Nadu, India

Abstract

Titanium dioxide nanoparticles (TiO₂-NPs) have a wide assortment of uses in numerous shopper items, including as food added substances, expanding the worry about the potential risks that TiO₂-NPs might posture to human wellbeing. These transformation pathways can have strong implications for the fate, transport, persistence, bioavailability and toxic-effects of the NPs. Besides, after inward breath or ingestion, TiO₂-NPs can arrive at a few organs, like the liver, cerebrum or lungs. Rate weight acquire (%WG) in treatment bunches showed critical contrasts from the basal eating regimen (T0) took care of gathering which underlines the significance of appropriate test conditions unfriendly impact on the development of fingerlings of *Labeo rohita*. This examination was directed to assess the impact of TiO₂-NPs on the development of fingerlings of *Labeo rohita*. TiO₂-NPs were utilized in various fixations (1 mg/kg, 10 mg/kg, 100 mg/kg and 1000 mg/kg) blended in fish basal feed for a time of 12 weeks. The death rate was recorded once the fish were accustomed, after TiO₂-NPs openness. The development long (cm) and weight (g) of the fish was estimated to assess the time and fixation subordinate development of the fingerlings. Hence, TiO₂-NPs in regular water bodies might influence the hydroponics business just as India fisheries.

Keywords: TiO₂-NPs, *Labeo rohita*, growth, final biomass, treatments

Introduction

Freshwater aquaculture in India contributes to over 95% of the total aquaculture production. The Indian significant carp, *Labeo rohita* is the most famous and overwhelming species refined in the Indian subcontinent. Titanium dioxide nanoparticles (TiO₂-NPs) are among the most mainstream fabricated and generally utilized nanoparticles [1]. They are delivered into the climate, influencing earthly and sea-going biological systems, with startling outcomes to creatures and human wellbeing. The current examination researches the interceded harmfulness forced to the freshwater fish species, zebrafish (*Danio rerio*) and the prussian carp (*Carassius gibelio*), and to the earthly land snail *Cornu aspersum*, after their openness to sublethal convergences of TiO₂-NPs [2]. Oxidative, proteolytic, genotoxic and apoptotic boundaries in fish liver and gills, just as on snail hemocytes were considered and the swimming execution was assessed to (a) gauge and recommend the most powerless creature, and (b) propose a typical battery of biomarkers as the most appropriate pointer for biomonitoring concentrates against TiO₂-NPs [3]. Our *in vivo* analyzes showed that NPs initiated adverse impacts on creature physiology and swimming conduct, while no broad example was seen in species and tissues responsiveness. By and large, TiO₂-NPs appeared to initiate a gathering of atoms that are normal for amphibian just as earthly creatures, suggesting the presence of a saved component. Titanium dioxide nanoparticles (TiO₂-NPs) are among the most well known made nanoparticles, progressively being joined into an assortment of mechanical and human items with respect to model paints, garments, beauty care products, drug, food and individual consideration items because of their high security, anticorrosion qualities and high photo catalytic action [4].

These metal oxides are delivered into the climate, influencing earthbound and amphibian environments, with unforeseen outcomes to living beings and people. They can enter the organic entities through different ways, where they are packed in organs, causing a few biologic responses and altering physicochemical properties in frameworks and tissues [5]. Various gatherings of creatures for example fish. As per the boundaries estimated each tissue reacted contrastingly against TiO₂-NPs openness. As our relationship results uncovered, it appears to be that liver in *C. gibelio* is more helpless than gills, when MDA and ubiquitin were thought of. In mouse liver cell the sign transducing occasions after TiO₂-NPs incorporate the cost like receptor initiation, that begins protein kinase phosphorylation and movement of NF-κB from the cytoplasm to the cores lastly prompts irritation [6, 7]. Additionally revealed liver as the most helpless organ to TiO₂-NPs openness, subsequent to looking at liver, gill and mind tissues of *Cyprinus carpio*. Broad n TiO₂ application and ubiquity of Cd might improve the shot at their co-openness to human. For instance, Cd-defiled food items contributed generously to add up to Cd admission by all inclusive community in numerous nations [8]. PGPB are additionally progressively being elevated as soil inoculants to work on rural efficiency, and as substitutes for compound composts [9]. With PGPB assuming a particularly essential part in soil fruitfulness, it becomes basic to assess the impact of TiO₂-NPs on these microscopic organisms. MeO-NPs are acquainted with the climate through various pathways, like waste expulsion (effluents, strong waste evacuation), deliberate applications for various purposes, and incidental delivery during creation and utilizations [10]. Among these courses, squander water release is the

fundamental pathway for the NPs delivery to the climate. Subsequently, different NPs are being recognized in different natural frameworks across the world. In the wake of being delivered, NPs can connect with various natural parts, go through changes and display distinctive conduct from their perfect structure ^[11]. The principle change measures are conglomeration/agglomeration, adsorption, affidavit, disintegration, redox responses, and association with macromolecules through different instruments, and these cycles are interlinked marvels of NPs in the climate. The cycles are influenced by practices of the NPs (for example size, morphology, surface charge), and states of the medium like IS, pH, temperature, and the presence of different parts (NOM, proteins, polysaccharides, extremists, surfactants) ^[12]. Be that as it may, fish exist in bound oceanic conditions (lakes, lakes, confines and so on), particularly the cultivated fishes in hostage condition are not equipped for evading them from every day or occasional changes in water temperature. Thusly, to sort out the impact of temperature changes on fish species, warm resistance and acclimation studies have seemed an impressive strategies for researchers. Acclimation temperature assumes a significant part in the warm resistance study. Acclimation can be characterized as the absolute capacity of fishes to adjust to short or long haul changes in their general climate. It has been seen that the life forms that show high physiological versatility more prominent capacity to acclimate to the new warm condition, can all the more likely make due in a warm variances or in a living space that go through unexpected warm changes ^[13]. Better Moreover, numerous microorganisms advance plant development in a roundabout way by forestalling multiplication of plant microbes. Antibacterial impacts of TiO₂-NPs have been generally concentrated on unadulterated societies of *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas aeruginosa*, and a couple clinical disconnects ^[14]. Since it is realized that TiO₂ has antibacterial action, it could affect sly affect microbes in soil. Microorganisms in soil play significant life-supporting jobs - reusing components, separating natural matter, advancing plant development, and so forth, hence adding to plant efficiency. An immediate advancement of plant development is achieved when these microorganisms utilize any at least one of the accompanying components - nitrogen obsession, phosphate solubilisation and union of phytohormones, siderophores, chemical 1-aminocyclopropane-1-carboxylate (ACC) de-aminase, hydrolytic catalysts, HCN or anti-toxins. Biofilm development is likewise a beneficial property in plant development advancing microbes (PGPB) as they can oppose abiotic stresses.

Material Methods

Fish collection

The fresh water Rohu (*Labeo rohita* Ham.) were collected from Saliyamangalam, Thanjavur District, Tamil Nadu, India.



Fig 1: (*Labeo rohita*)




Fig 2: Study area

Fish husbandry

Labeo rohita fingerlings, having a normal introductory load of 14.8 ± 0.15 g and an underlying length of 13 ± 0.11 cm, were solid fish of uniform sizes were chosen and given a KMnO₄ shower to eliminate any ectoparasites present. Preceding openness, fish were accustomed under 12 h light/dim routine in 100 L substantial tanks and took care of with basal eating regimen two times per day.

Diet formulation

The basal eating regimen (30% feed) and TiO₂ nanoparticles anatase (nano size < 25 nm, Sigma-Aldrich Corp, St. Louis, MO, USA) were taken from the neighborhood merchants. To acquire the test abstains from food, TiO₂ nanoparticles were added at 1, 10, 100 and 1000 mg/kg to the basal eating regimen, as per the exploratory convention. Molasses (7%) was utilized as a fastener to join the powder feed and structure a semisolid glue. This glue was gone through the mincing machine to shape noodles of 4 mm size/thickness. Noodles were cut into uniform little pellets, dried and saved in hermetically sealed containers for some time later ^[15].

Experimental design

The adjusted fish were separately gauged (g), estimated (cm) and conveyed in fifteen substantial tanks at 50 fish for each tank ($59 \times 34 \times 40$ cm) for all medicines three-fold replication. Medicines were recognized by the convergence of nanoparticles added to the feed as T0 (control), T1 (1 mg/kg), T2 (10 mg/kg), T3 (100 mg/kg) and T4 (1000 mg/kg). Fish were kept up with in a move through framework (stream rate 5.1–6 L hour₋₁) where water was kept up with at a moderate temperature ($29.4\text{--}30.1$ °C), was marginally antacid (pH = 7.4–7.6) and wealthy in disintegrated oxygen (11.1–11.5 mg/l) throughout the examination, in all medicines, as estimated with a HANNA multi-boundary test HI 9829. Fish were taken care of with test diet double a day (at 8 am and 8 pm) for thirty minutes to guarantee that all feed was eaten. Fish in the benchmark group were taken care of likewise with basal eating regimen till satiation. Uneaten feed and waste material were siphoned following the taking care of period. Fish were developed for a time of 12 weeks and any dead fish were taken out and recorded.

Fish sampling

Five fish from each tank (15 for every treatment) were inspected on day zero and later on fortnightly. Fish were euthanized by drenching them in aq. 3-aminobenzoic

corrosive ethyl ester (MS-222, Sigma-Aldrich Chemical, St Louis, MO; at 1.0 g/l), until the end of opercular development. The fingerlings were washed in clean water; paper dried gauged (g) and estimated for all out length (cm). Development boundaries like weight gain and explicit development rate (SGR) were determined by the accompanying equations a.) And b.). Physicochemical boundaries of the water like temperature, pH and broke down oxygen were estimated every day.

$$\text{Weight gain} = \frac{\text{Final weight} - \text{initial weight}}{\text{Initial weight}}$$

$$\text{Specific growth} = \frac{\text{rate Ln (Final weight)} - \text{Ln (Initial weight)}}{\text{Experimental periods in days}} \times 100$$

Statistical analysis

Comparison of various parameters was computed by using one way Analysis of Variance (ANOVA) and and Graph pad PRISM v.8.0. Correlation with repeated sampling at 95% level of significance. Data are represented as means ± SEM. Statistical significance of differences among different treatments was analyzed by Duncan’s Multiple Range test. All the statistical analysis was performed with software SPSS 24.0 version.

Result and Discussion

A higher mortality was observed in T3 and T4 as compared to T1 and T2, whereas no mortality was recorded in the T0 group, highlighting the lethal effect of TiO₂-NPs on the freshwater fish *Labeo rohita*. These outcomes are in close concurrence with the announced qualities for Oreochromis niloticus which showed 40%, half and 80% mortality at a nanoparticle convergence of 150, 175 and 200 mg/L individually, in 96 h. All things considered, the mortality was 100% at a nanoparticle convergence of 225 mg/L inside 24 h [16]. This species likely got away from raising homesteads around the Karun River because of reckless administration. Presentation of outlandish freshwater fishes is normal in hydroponics overall treating local biodiversity. Boundless use of titanium dioxide nanoparticles (n TiO₂) and pervasive cadmium (Cd) contamination might build their shot at concurrence in the common habitat. Toxicological data on co-openness of n TiO₂ and Cd in mammalian models is to a great extent lacking. Henceforth,

we considered the joined impacts of nTiO₂ and Cd in human liver (Hep G2) and bosom disease (MCF-7) cells [17]. We saw that n TiO₂ didn't create poisonousness to HepG2 andMCF-7 cells. Be that as it may, moderate convergence of Cd openness made cytotoxicity the two cells. Strangely, non-cytotoxic grouping of n TiO₂ adequately upgraded the oxidative pressure reaction of Cd demonstrated by supportive of oxidants age (responsive oxygen species, hydrogen peroxide, and lipid peroxidation) and cancer prevention agents consumption (glutathione level and glutathione reductase, superoxide dismutase, and catalase compounds). Particularly, fingerlings of *L. rohita* of gathering T4 showed an essentially (P < 0.05) higher death rate when contrasted and bunch T0. The mortality found in the current investigation may be because of different attributes of the TiO₂-NPs including the shape, size and sythesis. In the trials, the physicochemical boundaries of water like saltiness, temperature, broken down oxygen and pH were completely kept up with in the typical reach, appropriate for *Labeo rohita*, and the fish were taken care of similarly.

Table 1: proximate analysis of the experimental feed

S. no	Proximate analysis	Experimental feed %
1	Crude Protein	Min. 40%
2	Crude Fat	Min. 6.2%
3	Crude Fiber	Max. 6.9%
4	Moisture	Max. 10.5%
5	Ash	Max.8.7%

In the current examination, some conduct changes in the fish like imbalanced body, quake and vertical situating before death at high centralizations of TiO₂ nanoparticles were likewise recorded. These conduct changes are as per [18]. The body and social changes in zebra fish at high centralization of titanium dioxide nanoparticles for long haul openness were reported [19]. Likewise, high convergences of silver nanoparticles have been accounted for to prompt conduct changes in the fish and to deadly respiratory impacts in zebra fish, where surface breath and expanded operculum development were noticed [20]. The retarded growth in the fish species was probably outstanding to the production of reactive oxygen species (ROS) in the presence of TiO₂-NPs that hamper the cell membranes and cause oxidative stress related effects.

Table 2: Growth performance in fingerling of rohu (*Labeo rohita*), (WG-Weight Gain), (SGR -Specific Growth Rate) and (L- Length) fed with basal and experimental diets with different levels of nanoparticles given as Means ± S.E.

S. No	Parameters	Treatment	Initial Weight and weight Gains in n th week						
			Weight gain in n th week						
1			1 st	2 nd	4 th	6 th	8 th	10 th	12 th
2	WG	T0	5.27±0.01a	2.89±0.05a	1.07±0.07ab	2.07±0.08ab	1.77±0.07ab	3.20±0.05ab	1.69±0.05ab
		T1	7.37±0.04a	2.27±0.05a	5.57±0.05ab	6.7±0.002ab	8.5±0.005ab	9.57±0.05ab	9.87±0.05ab
		T2	5.21±0.03a	5.49±0.05a	7.37±0.05ab	9.37±0.02ab	11.7±0.07ab	13.0±0.07ab	14.0±0.02ab
		T3	4.17±0.05a	1.97±0.05a	0.39±0.05ab	4.27±0.10ab	3.27±0.05ab	2.07±0.02ab	1.39±0.05ab
		T4	4.04±0.05a	3.69±0.05a	2.87±0.05ab	1.7±0.040ab	1.07±0.02ab	0.70±0.05ab	0.56±0.05ab
S. No	Parameters	Treatment	Initial Weight and weight Gains in n th week						
			Weight gain in n th week						
1			1 st	2 nd	4 th	6 th	8 th	10 th	12 th
2	SGR	T0	1.29 ± 0.07a	1.77±0.02a	0.87 ± 0.05a	0.64±0.04a	1.07±0.18a	0.69±0.10a	0.57 ± 0.03a
		T1	1.0 ± 0.0a	1.75±0.05a	0.79 ± 0.03a	1.17±0.02a	0.67±0.05a	0.72±0.03a	0.72 ± 0.02a
		T2	1.59 ± 0.07a	1.00±0.07a	0.57 ± 0.05a	1.39±0.06a	1.04±0.02a	1.09±0.06a	0.51 ± 0.02a
		T3	0.97 ± 0.07a	0.68±0.02a	0.37± 0.02a	0.37±0.42a	0.67±0.04a	1.17±0.03a	0.9 ± 0.06a
		T4	0.37± 0.07a	0.39±0.06a	0.27± 0.02a	0.75±0.02a	1.17±0.03a	1.07±0.05a	1.07 ± 0.05a

S. NO	Parameters	Treatment	Initial Weight and weight Gains in n th week						
			Weight gain in n th week						
1			1 st	2 nd	4 th	6 th	8 th	10 th	12 th
	Length	T0	2.37 ± 0.07a	1.17±0.05a	0.79±0.05ab	0.28±0.05ab	0.59 ±0.05ab	0.39±0.05a	0.49±0.05ab
2		T1	3.27 ± 0.02a	1.79±0.05a	0.79±0.05ab	0.97 ±0.10ab	0.71 ±0.05ab	0.39±0.05ab	1.41±0.05ab
3		T2	1.97±0.02ab	1.36±0.05a	0.92±0.05ab	0.92 ±0.05ab	0.92 ±0.05ab	1.00±0.10ab	1.19±0.05ab
4		T3	1.97 ±0.05ab	1.36±0.05a	0.89±0.05ab	0.87 ±0.05ab	0.87±0.05ab	1.00±0.10ab	1.25±0.05ab
5		T4	0.67 ±0.02ab	0.47±0.05a	0.72±0.05ab	0.42 ±0.05ab	0.41 ±0.05ab	0.42±0.05ab	0.36±0.07ab

*Different letters indicate significant difference where ax < 0.05, ay < 0.01.

Protein is the most important organic material needed to construct and repair tissues and has an important role in providing energy for fish [21]. In fish, protein breakdown in order to provide energy and to deal with toxic effects of paraquat, intermediate metabolites, and a combination of

TiO₂ nanoparticles and paraquat (under darkness) may explain the significant decrease in total protein of plasma. Reduction of total protein has a significant effect on albumin and globulin levels in plasma of fish exposed to nano- TiO₂ and paraquat.




Fig 1: Growth performance in fingerling of rohu (*Labeo rohita*), A- (WG-Weight Gain), B- (SGR -Specific Growth Rate) and C- (L- Length) fed with basal and experimental diets with different levels of nanoparticles given as Means ± S.E

Table 3: Percentage weight gain (% WG in grams) and final biomass (FB) in different treatments (Means ± S.E).

S.No	Parameters -Treatment	Parameters -Treatment				
		Control	T1 (1mg/kg)	T2 (10 mg/kg)	T3 (100 mg/kg)	T4 (1000 mg/kg)
1	%WG	91.47 ± 1.42a	129.81 ± 0.20ax	147.45 ± 0.60ax	100.97 ± 0.21ax	85.38 ± 0.77ay
2	Final biomass (g)	29.7 ± 1.27a	35.27 ± 1.07ax	37.0 ± 0.5ax	29.17 ± 0.99ay	27.5 ± 0.67ay

Since monitor the effects of pollutants on varied organisms is difficult, owing to their broad range of responses to a known pollutant, it is critical to estimate the response of a group of animal and single animal model. To our knowledge, this is the first study that compares the effect of

the widely used TiO₂-NPs on the responsiveness of tissues of three animal models, of both aquatic and terrestrial environments, in order to suggest reliable biomarkers and bio indicators against NPs [22]. Our *in vivo* analyzes exhibited that NPs actuated impeding impacts on creature physiology and swimming conduct. By and large, TiO₂-NPs appear to initiate a gathering of atoms that are normal for sea-going just as earthbound creatures, inferring the presence of a saved component [23]. It appears to be that after openness to TiO₂-NPs, a typical instrument is actuated that includes the incitement of resistant framework with the creation of ROS, harm of lysosomal layer, protein carbonylation, lipid peroxidation, DNA harm, following proteolysis by ubiquitin lastly apoptosis. Consequently, the concurrent utilization of the last biomarkers could be recommended as a dependable multi boundary approach for

bio monitoring of amphibian and earthy environments against TiO₂-NPs [24]. At long last, the conditions applied in the current investigation and the centralizations of TiO₂-NPs tried, appearing to prompt oxidative damage to creature cells, without influencing their endurance. Further investigations of TiO₂-NPs initiated harmfulness on creature biochemical and physiological reactions are fundamental, to guarantee their protected use and spillage to the climate. Contaminations including nano-sized materials released into water bodies influence oceanic environments by means of their take-up by the amphibian living beings and their collection in creature tissues after some time prompting resulting gastrointestinal harmfulness. Who researched two marine fish, red drum (*Sciaenops ocellatus*) and marine medaka (*Oryzias melastigma*) which were presented present moment to nano-ZnO nano particles. Constructive outcomes were recorded at 80 mg/kg nano-ZnOs, while high dietary expansion of ZnO-NPs (300 mg/kg) prompted hindered development in medaka (stomach-less fish). Generally speaking, it should be noticed that the current investigation shows that the openness of fingerlings of *L. rohita* to TiO₂-NPs at convergences of 100 mg/kg feed or more contrarily influences the development pace of the fish fundamentally. Death rates as of now increment at lower focuses yet in addition become huge at centralizations of 100 mg TiO₂-NP/kg feed or more.

Summary and Conclusion

TiO₂-NPs help to advance the development of fish as far as body weight, rate weight gain and SGR. Long haul openness of TiO₂nanoparticles at high focuses was demonstrated to be poisonous and surprisingly deadly for youthful freshwater fish. Sea-going (water and dregs) and earthy compartments are anticipated to be the objective of the delivered MeO-NPs. In these compartments, the particles are exposed to different unique cycles like physical, compound and organic cycles, and go through changes which drive them away from their perfect state [25]. These change pathways can have solid ramifications for the destiny, transport, determination, bioavailability and harmful impacts of the NPs. In this basic survey, we give the condition of-the-information on the change cycles and bioavailability of MeO-NPs in the climate, which is the subject important to specialists. Show that NP take-up relies upon the focus, time and cell type. Our investigation additionally builds up the significance of cell model and testing convention decision for poisonousness appraisal. Taking into account that TiO₂-NPs were electively disguised by all cell lines, yet no genotoxicity was related, current information recommend that TiO₂-NPs don't a chromosomal honesty. The methodology here introduced might be valuable in the genotoxicity appraisal of different kinds of NPs, basically reference materials in cell lines of various beginning and delegates of NPs target organs. File of Preponderance (IOP) uncovered that green growth, alongside protozoans, rotifers and cladocerans were the most favored food things in the more modest size gatherings and denoted a decay as the size of the fish expanded. In the high level stages, macrophytes, garbage and sand/mud were found to frame a significant portion of food. In this way rohu has been classified as omnivore-planktophage in beginning phases of life and herbi-omnivore in grown-up stages. As found in our investigation, powerlessness of Gram-negative microorganisms even in bunches with comparative qualities

was totally different. For instance, among the biofilm formers, *Azotobacter* and *Rhizobium* were not as safe as *Klebsiella*, *Pseudomonas* and *Serratia*. The idea of LPS alterations and lipids in the external layer perhaps answerable for the noticed contrasts in the vulnerability of Gram-negative microscopic organisms. There is a need to test a more extensive assortment of microbes, essentially of various genera, utilizing normalized systems to assess impacts of TiO₂-NPs. Plant development elevating microorganisms having a place with three gatherings, *viz.*, nitrogen fixers, phosphate solubilizers and biofilm formers, were altogether hindered in direct contact with TiO₂-NPs. The restraint showed a stage reaction which was disposed of within the sight of rich development medium. The nanoparticles give off an impression of being in hibitorier to Gram-positive microscopic organisms recommending that bacterial surface might play a significant part to play in harmfulness of TiO₂-NPs.

Acknowledgement

The author is also very grateful to the President of our College Management Committee and has been unanimously elected as the Secretary and Correspondent and the Principal of A.V.V.M. Sri Pushpam College (Autonomous), Poondi-613 503, Thanjavur (Dt), for providing the excellent infrastructure and necessary facilities to carry out my research work successfully.

Conflict of Interest

The authors have declared that there is no conflict of interest.

Source/S of Funding

No source of funding.

References

- Ahamed M, Akhtar MJ, Alaizeri ZAM, Alhadlaq HA.. TiO₂ nanoparticles potentiated the cytotoxicity, oxidative stress and apoptosis response of cadmium in two different human cells. *Environmental Science and Pollution*,2020;27:10425-10435.
- Areecheewakul S, Adamcakova-Dodd A, Givens BE, Steines BR, Wang Y, Meyerholz DK *et al.* Toxicity assessment of metal oxide nanomaterials using *in vitro* screening and murine acute inhalation studies. *Nano Impact*,2020;18:100214.
- Abbas F, Qureshi NA, Khan N, Ashraf M, Iqbal KJ. Study on the digestibility and growth potential of artificial feed in *Catla catla*, *Cirrhinus mrigala* and *Labeo rohita*. *The Journal of Animal and Plant Sciences*,2019;29:695-702.
- Baranowska-Wójcik E, Szwajgier D, Oleszczuk P, Winiarska-Mieczan A. Effects of titanium dioxide nanoparticles exposure on human health-a review. *Biological Trace Element Research*,2020;193:118-129.
- Rahman MM, Alam MM, Asiri AM, Islam MA. Fabrication of selective chemical sensor with ternary ZnO/SnO₂/Yb₂O₃ nanoparticles. *Talanta*, 2017;170:215e223.
- Bobori D, Dimitriadi A, Karasiali S, Tsoumaki-Tsouroufli P, Mastora M, Kastrinaki G *et al.* Common mechanisms activated in the tissues of aquatic and terrestrial animal models after TiO₂ nanoparticles exposure. *Environment International*,2020;138:105611.

7. Brandão F, Fernández-Bertólez N, Rosário F, Bessa MJ *et al.* Genotoxicity of TiO₂ nanoparticles in four different human cell lines (A549, HEPG2, A172 and SHSY5Y). *Nanomaterials*, 2020;10(3):412. <https://doi.org/10.3390/nano10030412>.
8. Romero-Freire A, Lofts S, Martin Peinado FJ, van Gestel CA. Effects of aging and soil properties on zinc oxide nanoparticle availability and its ecotoxicological effects to the earthworm *Eisenia andrei*. *Environ. Toxicol. Chem*,2017;36:137e146.
9. Cai J, Zhuo X, Yan X, Lucente D, Lagana C. Top 10 species groups in global aquaculture, Food and Agricultural Organization of the United Nations, 2019. <http://www.fao.org/3/ca5224en/ca5224en.pdf> (accessed), 2020, 4.
10. Chavan S, Sarangdhar V, Nandanathangam V. Toxicological effects of TiO₂ nanoparticles on plant growth promoting soil bacteria. *Emerging Contaminants*,2020;6:87-92.
11. Saba MK, Amini R. Nano-ZnO/carboxymethyl cellulose-based active coating impact on ready-to-use pomegranate during cold storage. *Food Chem*, 2017;232:721e726.
12. Dong LL, Wang HX, Ding T, Li W, Zhang G. Effects of TiO₂ nanoparticles on the life-table parameters, antioxidant indices, and swimming speed of the freshwater rotifer *Brachionus calyciflorus*. *Journal of Experimental Zoology*,2020;333(4):230-239.
13. Du J, Xu S, Zhou Q, Li H, Fu L, Tang JH, Jin MQ. The ecotoxicology of titanium dioxide nanoparticles, an important engineering nanomaterial. *Environmental Toxicology and Chemistry*,2019;101:165-189.
14. Khan I, Saeed K, Khan I. Nanoparticles: properties, applications and toxicities. *Arabian Journal of Chemistry*,2019;12(7):908-931.
15. Faiz H, Zuberi A, Nazir S, Rauf M, Younus N. Zinc oxide, zinc sulfate and zinc oxide nanoparticles as source of dietary zinc: Comparative effects on growth and hematological indices of juvenile grass carp (*Ctenopharyngodon idella*). *International Journal of Agriculture and Biological*,2015;17(3):568-574.
16. Kumar S, Sahu NP, Deo AD, Ranjan A. Feeding de-oiled rice bran based diet with varying level of protein and lipid: Effect on physiological responses of *Labeo rohita*. *Aquaculture*,2019;498:454-463.
17. Sendra M, Yeste MP, Gatica JM, Moreno-Garrido I, Blasco J. Homoagglomeration and heteroagglomeration of TiO₂, in nanoparticle and bulk form, onto freshwater and marine microalgae. *Sci. Total Environ*, 2017a;592:403e411.
18. Matet A, Behar-Cohen F, Cassoux N, Declèves X, Cisternino S. Retinal and choroidal cancers: Blood-retinal barriers considerations in ocular chemotherapy. In: *Drug efflux pumps in cancer resistance pathways: From molecular recognition and characterization to possible inhibition strategies in chemotherapy*. Elsevier, 2020, 303-335
19. Peng YH, Tsai YC, Hsiung CE, Lin YH, Shih YH. Influence of water chemistry on the environmental behaviors of commercial ZnO nanoparticles in various water and wastewater samples. *J. Hazard. Mater*, 2017;322:348e356.
20. Yu Q, Wang H, Peng Q, Li Y, Liu Z, Li M. Different toxicity of anatase and rutile TiO₂ nanoparticles on macrophages: involvement of difference in affinity to proteins and phospholipids. *J. Hazard. Mater*, 2017;335:125e134.
21. De Loid GM, Cohen JM, Pyrgiotakis G, Demokritou P. Preparation, characterization, and *in vitro* dosimetry of dispersed, engineered nanomaterials. *Nature protocols*, 2017a;12(2):355-371. <https://doi.org/10.1038/nprot.2016.172>. (Epub 2017/01/20, PubMed PMID: 28102836; PMCID: PMC5857388).
22. Vidya PV, Chitra KC. Assessment of acute toxicity (LC50-96 h) of aluminium oxide, silicon dioxide and titanium dioxide nanoparticles on the freshwater fish, *Oreochromis mossambicus* (Peters, 1852). *International Journal of Fisheries and Aquatic Studies*,2017;5(1):327-332.
23. Zande M, Undas AK, Kramer E, Monopoli MP, Peters RJ, Garry D *et al.* Different responses of Caco-2 and MCF-7 cells to silver nanoparticles are based on highly similar mechanisms of action. *Nanotoxicology*, 2016;10:1431-1441.
24. Zeumer R, Galhano V, Monteiro M, Kuehr S, Knopf B, Meisterjahn B *et al.* Chronic effects of wastewater-borne silver and titanium dioxide nanoparticles on the rainbow trout (*Oncorhynchus mykiss*). *Science of the Total Environment*,2020;723:137974.
25. Picchietti S, Bernini C, Stocchi V, Taddei AR, Meschini R, Fausto AM *et al.* Engineered nanoparticles of titanium dioxide (TiO₂): uptake and biological effects in a sea bass cell line. *Fish. Shellfish Immunol*,2017;63:53e67.